



Measuring Uranium Concentration in Amara Governorate, Southern Iraq in Two Different Ways

Hawraa J. Jasim, Thaer M. Salman &  Mostafa Ahmed Algrifi*

Department of Physics, College of Education for Pure Science, University of Basrah, Basrah, Iraq

*Corresponding author: Mostafa Ahmed Algrifi | e-mail: mostafajawad88@gmail.com

ARTICLE INFO

Article history:

Received on: April 10, 2025

Revised on: June 19, 2025

Accepted on: June 26, 2025

Published on: July 01, 2025

Keywords:

CR-39 detector

ICP-MS

Neutron activation technique

Soil

SSNTDs

Uranium concentration

ABSTRACT

The current study uses ICP-MS (inductively coupled plasma mass spectrometry) and SSNTDs (solid-state nuclear track detectors) to measure uranium concentrations in 25 surface soil samples from selected locations in Omara governorate in Southern Iraq. In the ICP-MS technique, uranium concentrations ranged from 0.66 ppm to 1.6 ppm. In the SSNTDs technique, uranium concentrations ranged from 0.81 ± 0.31 ppm to 1.95 ± 0.50 ppm. The results obtained from the soil samples were within the permissible limits recommended by the UNSCEAR Committee of 11.7 ppm.

Copyright © 2025 *Biomedicine and Chemical Sciences*. Published by *International Research and Publishing Academy* – Pakistan, Co-published by *Al-Furat Al-Awsat Technical University* – Iraq. This is an open access article licensed under CC BY: (<https://creativecommons.org/licenses/by/4.0>)

INTRODUCTION

Uranium is a silvery, shiny metal with a long half-life as a natural radionuclide. Uranium is the heaviest element in nature. It is one of the most significant environmental dangers due to its radioactivity. Uranium and its compounds are very toxic and threaten human health and the environment (Algrifi & Salman, 2023). Uranium is a common element that may exist as a solid, liquid, or gas. It may be contained in food, water, soil, rocks, natural materials, and the environment. When uranium combines with other elements, it easily creates uranium oxide, silicates, hydroxides, and carbonates (Algrifi & Salman, 2023). Their solubility determines the physiological action of uranium compounds. The chemical toxicity of soluble uranium is regulated, but the radiological properties of insoluble (less soluble) uranium are not. However, because of its slow absorption via the lungs and long retention time in human tissues, it will significantly

harm internal organs through radiological damage (cancer mortality risk) rather than a chemical hazard to the kidneys (Al-Hamzawi et al., 2015). Uranium may enter the human body through a variety of pathways. It enters the body either by breathing uranium-containing dust particles, drinking uranium-contaminated water, or indirectly through the food chain from the fertile soil layer (Chan & Hieftje, 2013).

This study aims to examine the uranium content in selected soil samples taken from various residential, industrial, and agricultural sectors in the Um Qasr region in southern Iraq using ICP-MS technology and neutron activation technique for nuclear pathway detectors CR-39. This study was conducted in the Um Qasr region due to the lack of previous research and the establishment of a database of uranium concentrations in soil samples. The study site was chosen because of its importance in terms of the use of many weapons in it, and these two techniques were chosen to obtain the best results.

How to Cite:

Jasim, H. J., Salman, T. M., & Algrifi, M. A. (2025). Measuring Uranium Concentration in Amara Governorate, Southern Iraq in Two Different Ways. *Biomedicine and Chemical Sciences*, 4(3), 82–87. <https://doi.org/10.5281/zenodo.16651823>

Publisher's Note:

International Research and Publishing Academy (iRAPA) stands neutral with regard to jurisdictional claims in the published maps and institutional affiliations.

Copyright:

© 2025 | *Biomedicine and Chemical Sciences* published by *International Research and Publishing Academy* (iRAPA)



This is an Open Access article published under the Creative Commons Attribution 4.0 International (CC BY 4.0) (<https://creativecommons.org/licenses/by/4.0>)

Creative Commons Attribution (CC BY): lets others distribute and copy the article, to create extracts, abstracts, and other revised versions, adaptations or derivative works of or from an article (such as a translation), to include in a collective work (such as an anthology), to text or data mine the article, even for commercial purposes, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, and do not modify the article in such a way as to damage the author's honour or reputation.

Toxicity of Uranium

The chemical and radiological consequences of uranium and its derivatives are hazardous. The renal and respiratory effects of uranium exposure in humans and animals are usually attributed to its chemical properties. In contrast, hypothetically, probable excess malignancies are generally induced by its radiation properties (Cléro et al., 2019).

Chemical Toxicity of Uranium

The chemical toxicity of an elemental compound is related to the substance's interaction with the biochemical processes of the human body. Some of these interactions may be beneficial or even required, whereas others may be detrimental (Craig, 2001). Uranium's toxicity is a crucial aspect. Its chemical and radioactive characteristics determine the harmful effects of uranium exposure. The toxicity of uranium compounds is directly related to their solubility; the more soluble a uranium compound is, the more poisonous it is; soluble uranium compounds are harmful when breathed in or eaten. UF_6 , $UO_2(NO_3)_2$, UO_2Cl_2 , UO_2F_2 , and carbonates (Dockery et al., 1993) are the most soluble compounds of uranium. When uranium hexafluoride (UF_6) is emitted into the air, it quickly interacts with water vapor, forming uranium oxyfluoride and hydrogen fluoride. Because the chemical reaction produces heat, a plume of these chemical compounds may become buoyant and ascend. The substances produced by the result cause three toxic consequences:

- The uranyl complex contains uranium, a heavy metal toxin that can harm the kidneys
- Hydrogen fluoride is an acid that, when concentrated, may inflict acid burns on the skin or lungs
- If excessive amounts of fluorides (uranium oxyfluoride and hydrogen fluoride) are consumed, fluoride poisoning can occur

Radiological Toxicity

Dosage of radiation is critical in radiation protection (ionizing radiation energy absorbed per unit mass by a target organ or tissue). The ability of various radionuclides' ionizing radiation to infiltrate materials varies depending on the kind of radiation and its energy. External and internal radionuclide exposures are also possible, and their relative importance varies depending on the type of radiation and radionuclides involved. Because of their massive size and charge, alpha particles quickly lose their kinetic energy and have little penetrating power; hence, the health

repercussions of external exposure are limited to skin contact. As a consequence, assuming no uranium redistribution or sorption into the epidermis, the penetrating range for an alpha particle from ^{238}U in soft tissue is approximately 28 μ m. Alpha particles cannot penetrate the outer layer of skin when discharged outside the body and do not represent a hazard. Beta particles can penetrate the outer layers and deliver a localized dose when they come into contact with the skin. Gamma radiation can penetrate further into the body and deposit energy in interior organs, depending on the strength of the gamma radiation emitted. The precise mechanisms by which radiation interacts with biological materials are still being investigated. However, the energy of radiation may injure cells by causing changes in deoxyribonucleic acid (DNA). This physiologically necessary molecule governs cell structure and function and is found mainly in cell nuclei, and ionizing radiation has been shown to damage individual chromosomes. DNA damage or fragmentation is the most prevalent negative effect of low-level ionizing radiation exposure. Viable cells heal the damage; errors might produce gene mutations or chromosomal abnormalities. The risk of lung cancer has been connected to alpha particle radiation from ^{222}Rn and its decay products, which occur in uranium mines due to the dissolution of ^{238}U . Inhaled particle irradiation of the lungs is principally responsible for the enhanced toxicity of insoluble compounds. The ability of various radionuclides' ionizing radiation to infiltrate materials varies depending on the radiation emitted and its energy (Durrani & Bull, 2013; Salman & Algrifi, 2022).

MATERIALS & METHODS

Collection of Soil Samples

In this study, 25 soil samples were obtained from 25 separate places in southern Iraq's Omara governorate, one from each. Soil samples were taken at the depth inside the soil of 15cm (at one depth to obtain soil free of impurities). The research areas, which included residential, agricultural, and industrial zones, are depicted in Table 1. After cleaning and removing stones, pebbles, and root sections from 12 soil samples, they were dried in an oven at 70°C for a few hours before being powdered and using a special sieve with a diameter of 75 μ m for the sieve holes (Salman & Algrifi, 2022).

Analytical Determination of Uranium

The presence of uranium in the soil is noteworthy due to its extreme chemical toxicity. This section

summarizes several analytical techniques used in this research to determine uranium in different soil samples. Among the most used methods are:

Inductively Coupled Plasma Techniques /ICP-MS

Inductively Coupled Plasma Techniques have the potential to be highly effective techniques for identifying and studying trace and ultra-trace components. It's an intriguing alternate way of determining uranium. This approach offers various advantages, including a rapid analysis time and low detection limits. Low sample consumption with little spectrum interferences (Salman & Algrifi, 2022b). ICP-MS has been the technique of choice in many analytical laboratories in recent years for providing the accurate and exact measurements required for today's demanding applications and the required lower limits of detection. Mass spectrometry has high sensitivity as well as accuracy. In ICP-MS, Argon gas is used to create plasma or gas containing ions, electrons, and neutral particles, which are then used to atomize and ionize the elements in the sample matrix. These ions are then passed via a succession of apertures (cones) into a high vacuum mass analyzer, where the element isotopes are identified based on their mass-to-charge ratio. The strength of a particular peak in the mass spectrum is related to the quantity of the elemental isotope present in the original sample (Salman & Algrifi, 2022b).

A high-temperature ICP (Inductively Coupled Plasma) source is combined with a mass spectrometer in an ICP-MS. The atoms of the elements in the sample are converted to ions by the ICP source. The mass spectrometer then separates and detects these ions. Ions created in the plasma (at atmospheric pressure) flow via a sampler and skimmer cone assembly to a highly evacuated MS region in ICP-MS. Ions are focused on an ion optic system, which monitors the mass-to-charge ratios of the ions. The common quadrupole MS works as a mass filter, allowing only a specific mass-to-charge ratio of ions to get through to the detector. The ions moving through the MS are deflected to an ion detector, which transforms the ionic energy into electric energy, allowing the analytic concentration to be measured (Todorov & Ilieva, 2006).

Neutron Induced Fission Track Technique

Many researchers have used the neutron-induced approach to determine uranium in solid materials.

The ^{235}U (n,f) reaction is used in this approach. The primary idea behind this approach is to employ the ^{235}U (n,f) reaction, which has a fission cross-section of 4.2 neutrons. The soil samples were dried in an electric oven at 70°C for 6 hours, after which they were ground with a grinder. (0.5 g) powder soil was combined with (0.1 g) methylcellulose as a binder. A manual piston with a diameter of 1 cm and a thickness of 1.5 mm was used to crush the mixture into a pellet. A CR-39 track detector with a size of (1.5×1.5 cm) was coated on both sides of the pellets. The pellets were subsequently irradiated in a paraffin wax dish for 7 days at a distance of 5 cm from the neutron source (^{241}Am - ^9Be) source with a flux of (2.3×10^5 n/cm².s) to cause latent damage on the detector due to the ^{235}U reaction (n,f). After the irradiation technique, the detectors are etched in a (6.25N) NaOH solution at a temperature of 70 °C for (6 h). The detectors are removed at the end of the etching process, washed in distilled water, and dried. The optical microscope was then used to count the number of alpha tracks (US Department of Health & Human Services, 1990). As a result, if an unknown soil sample and a known uranium standard are both irradiated and brought into contact with the detector at the same time in the reactor with the same integrated flux of thermal neutrons, the uranium content of the unknown sample is calculated after chemical etching by calculating and comparing the total number of thermal neutrons in the reactor. Observed fission routes arising in detectors owing to uranium in samples and expected result (n, f) nuclear reaction in the reactor. This technique makes use of SSNTDs. Solid-state nuclear trajectory detectors have become essential instruments in many fields of research and technology, including neutron dosimetry, gamma ray detection, cosmic rays, heavy ions, and nuclear physics. SSNTDs have been adjusted and evaluated in several laboratories to increase detection sensitivity, charge accuracy, and power (Zarkadas et al., 2001).

RESULTS & DISCUSSION

Table 1 presents uranium concentrations for soil samples measured by ICP/ MS. The uranium concentrations range from 0.66 ppm to 1.6 ppm. The maximum uranium concentration of 1.6 ppm was in station number S9. The minimum value of 0.66 ppm was observed in station number S24 as shown in Figure 1.

Table 1
The uranium concentration of soil sample in Omara by ICP -MS

Sites numbers	Sites	Uranium Concentration in ppm
S1	Alwadia1	0.8
S2	Alwadia2	0.96
S3	Alwadia3	0.79
S4	Almaymuna1	0.95
S5	Almaymuna2	1.05
S6	Almaymuna3	1.22
S7	Almaymuna4	1.31
S8	Almaymuna5	1.55
S9	Almaymuna6	1.6
S10	Almaymuna7	1.1
S11	Almaymuna8	0.92
S12	Alkahla1	0.89
S13	Alkahla2	0.99
S14	Alkahla3	0.93
S15	Alkahla4	0.84
S16	Alkahla5	0.9
S17	Alkahla6	1.33
S18	Alkahla7	0.98
S19	Aluzair1	1.49
S20	Aluzair2	1.21
S21	Aluzair3	1.15
S22	Aluzair4	1.37
S23	Aluzair5	1.29
S24	Aluzair6	0.66
S25	Aluzair7	0.74

Table 2 presents uranium concentrations for soil samples in Um Qasr district measured using the irradiation method (the CR-39 detector). The uranium concentrations for soil range from 1.95 ± 0.50 ppm to 0.81 ± 0.31 ppm. The highest concentration for

uranium, 1.95 ± 0.50 ppm, was observed, as shown in Figure 1, which indicates that this site is higher than other regions, whereas the lowest value is 0.81 ± 0.31 ppm.

Table 2
Uranium Concentration and specific activities in Omara in soil by SSNTDs

Sites numbers	Sites	Tracks density(tracks/mm ²)	Uranium Concentration in ppm
S1	Alwadia1	460.54	1.12±0.36
S2	Alwadia2	493.44	1.2±0.39
S3	Alwadia3	435.87	1.06±0.35
S4	Almaymunah1	514	1.25±0.54
S5	Almaymunah2	625.02	1.52±0.29
S6	Almaymunah3	694.92	1.69±0.52
S7	Almaymunah4	777.16	1.89±0.54
S8	Almaymunah5	801.84	1.95±0.50
S9	Almaymunah6	786.62	1.913±0.40
S10	Almaymunah7	620.91	1.51±0.39
S11	Almaymunah8	555.12	1.35±0.56
S12	Alkahla1	542.78	1.32±0.49
S13	Alkahla2	518.11	1.26±0.41
S14	Alkahla3	501.66	1.22±0.40
S15	Alkahla4	468.76	1.14±0.33
S16	Alkahla5	534.56	1.3±0.63
S17	Alkahla6	731.93	1.78±0.55
S18	Alkahla7	522.22	1.27±0.54
S19	Aluzair1	736.04	1.79±0.53
S20	Aluzair2	604.46	1.47±0.38
S21	Aluzair3	559.23	1.36±0.70
S22	Aluzair4	699.04	1.7±0.52
S23	Aluzair5	633.24	1.52±0.41
S24	Aluzair6	333.07	0.81±0.31
S25	Aluzair7	382.41	0.93±0.41



Fig. 1. Uranium concentration in soil samples from Omara

CONCLUSION

Uranium content in soil samples was determined using ICP-MS and SSNTDs. This study's findings revealed that uranium concentration increases in industrial regions more than in residential areas because the industrial areas contain repair workshops and the disposal of industrial wastes in them, thus

increasing the concentration of this element in these areas. However, the results were within acceptable levels and not cause for concern (the levels set by the World Health Organization).

Competing Interest

The authors had no competing interests.

REFERENCES

- [1] Algrifi, M. A., & Salman, T. M. (2023). Measuring uranium concentration in Um Qasr District, Southern Iraq, in two different ways. *Applied Radiation and Isotopes*, 192, 110595. <https://doi.org/10.1016/j.apradiso.2022.110595>
- [2] Al-Hamzawi, A. A., Jaafar, M. S., & Tawfiq, N. F. (2015). Concentration of uranium in human cancerous tissues of Southern Iraqi patients using fission track analysis. *Journal of Radioanalytical and Nuclear Chemistry*, 303(3), 1703-1709. <https://doi.org/10.1007/s10967-014-3682-0>
- [3] Chan, G. C. Y., & Hieftje, G. M. (2013). Spatial emission profiles for flagging matrix interferences in axial-viewing inductively coupled plasma-atomic emission spectrometry: 1. Profile characteristics and flagging efficiency. *Analytical Chemistry*, 85(1), 50-57. <https://doi.org/10.1021/ac302095w>
- [4] Cléro, E., Vaillant, L., Hamada, N., Zhang, W., Preston, D., Laurier, D., & Ban, N. (2019). History of radiation detriment and its calculation methodology used in ICRP Publication 103. *Journal of Radiological Protection*, 39(3), R19. <https://doi.org/10.1088/1361-6498/ab294a>
- [5] Craig, D. K. (2001). Chemical and radiological toxicity of uranium and its compounds. *Washington, DC, Westinghouse Savannah River Company*.
- [6] Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., ... & Speizer, F. E. (1993). An association between air pollution and mortality in six US cities. *New England Journal of Medicine*, 329(24), 1753-1759. <https://doi.org/10.1056/NEJM199312093292401>
- [7] Durrani, S. A., & Bull, R. K. (2013). *Solid state nuclear track detection: principles, methods and applications* (Vol. 111). Elsevier.
- [8] Salman, T. M., & Algrifi, M. A. (2022). Measurement of uranium concentration in Basrah soils using the CR-39 detector. *Biomedicine and Chemical Sciences*, 1(3), 164-167. <https://doi.org/10.48112/bcs.v1i3.174>

- [9] Salman, T. M., & Algrifi, M. A. (2022b). Using the ICP-MS method, the concentration of uranium in soil samples from the Northern Basrah governorate was determined. *Al-Bahir Journal for Engineering and Pure Sciences*, 1(1), 1. <https://doi.org/10.55810/2313-0083.1002>
- [10] Todorov, P. T., & Ilieva, E. N. (2006). Contamination with uranium from natural and antropological sources. *Romanian Journal of Physics*, 51(1/2), 27.
- [11] US Department of Health & Human Services. (1990). *Toxicological profile for uranium*.
- [12] Zarkadas, C., Karydas, A. G., & Paradellis, T. (2001). Determination of uranium in human urine by total reflection X-ray fluorescence. *Spectrochimica Acta Part B: Atomic Spectroscopy*, 56(12), 2505-2511. [https://doi.org/10.1016/S0584-8547\(01\)00348-2](https://doi.org/10.1016/S0584-8547(01)00348-2)